

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 March 2001 (01.03.2001)

PCT

(10) International Publication Number
WO 01/14537 A1

(51) International Patent Classification⁷: C12N 15/00,
15/63, A01K 67/00, 67/027

(21) International Application Number: PCT/US00/22433

(22) International Filing Date: 16 August 2000 (16.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/377,066 19 August 1999 (19.08.1999) US

(71) Applicant: THE UNITED STATES OF AMERICA,
as represented by THE SECRETARY OF AGRICULTURE [US/US]; 1400 Independence Avenue S.W.,
Washington, DC 20250-0302 (US).

(72) Inventor: HANDLER, Alfred, M.; 6825 Northwest 52nd
Terrace, Gainesville, FL 32653 (US).

(74) Agents: PENDORF, Stephan, A. et al.; Pendorf & Cutliff,
P.O. Box 20445, Tampa, FL 33622-0445 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/14537 A1

(54) Title: PIGGYBAC TRANSFORMATION SYSTEM

(57) Abstract: The present invention is directed to a transformation system for making transgenic organisms that includes a vector containing a modified *piggyBac* transposon into which is inserted an enhanced green fluorescent protein gene linked to a polyubiquitin promoter sequence and a nuclear localizing sequence; and a helper transposase vector that includes an *hsp70* promoter sequence upstream of the putative *piggyBac* promoter that increases the transformation frequency of this system.

PIGGYBAC TRANSFORMATION SYSTEM

BACKGROUND OF INVENTION

Field of the Invention

This invention relates to a transformation system that includes a gene transfer vector containing a modified piggyBac transposon (pB) and having the insertion of a marker construct containing an enhanced green fluorescent protein gene (EGFP) linked to a polyubiquitin promoter gene and a nuclear localizing sequence. The invention further relates to a helper vector containing a heat shock protein gene and to methods for using this system to transform eukaryotic cells as well as transgenic organisms produced using the system, especially insect cells and insects, respectively.

Description of the Related Art

The piggyBac transposable element from the cabbage looper moth, *Trichoplusia ni* (Cary et al., Virology, Volume 161, 8-17, 1989) has been shown to be an effective gene-transfer vector in the Mediterranean fruit fly, *Ceratitidis capitata* (Handler et al., Proc. Natl. Acad. Sci. USA, Volume 95, 7520-7525, 1998). Use of an unmodified transposase helper under piggyBac promoter regulation indicates that piggyBac retains autonomous function in the medfly, since transcriptional regulation was maintained, as well as enzymatic activity. This observation was unique since all other successful insect germline transformations had been limited to dipteran species using vectors isolated from the same or another dipteran. The initial transformation of medfly (Loukeris et al., Science, Volume 270, 2002-2005, 1995) used the Minos vector from *Drosophila hydei* (Franz & Savakis, Nucl. Acids Res., Volume 19, 6646, 1991), and *Aedes aegypti* has been transformed from Hermes (Jasinskiene et al., Proc. Natl. Acad. Sci. USA, Volume 95, 3743-3747, 1998) from *Musca domestica* (Warren et al., Genet. Res. Camb., Volume 64, 87-97, 1994) and mariner (Coates et al., Proc. Natl. Acad. Sci. USA, Volume 95, 3748-3751, 1998) from *Drosophila mauritiana* (Jacobson et al., Proc. Natl. Acad. Sci.

USA, Volume 83, 8684-8688, 1986). *Drosophila melanogaster* has been transformed as well by *Hermes* (O=Brochta et al., Insect Biochem. Molec. Biol., Volume 26, 739-753, 1996) *mariner* (Lidholm et al., Genetics, Volume 134, 859-868, 1993), *Minos* (Franz et al., Proc. Natl. Acad. Sci. USA, Volume 91, 4746-4750, 1994) and by the *P* and *hobo* transposons originally discovered in its own genome (Rubin and Spradling, 1989; Blackman et al., EMBO J., Volume 8, 211-217, 1989). *Drosophila virilis* also has been transformed by *hobo* (Lozovskaya et al., Genetics, Volume 143, 365-374, 1995; Gomez & Handler, Insect Mol. Biol., Volume 6, 1-8, 1997) and *mariner* (Lohe et al., Genetics, Volume 143, 365-374, 1996). While the restriction to dipteran vectors is due in part to the limited number of transposon systems available from non-dipteran species, phylogenetic limitations on transposon function is not unexpected considering the deleterious effects functional transposons may have on a host genome. This is, indeed, reflected by the high level of regulation placed on transposon movement among species, among strains within a host species, and even among cell types within an organism (Berg & Howe, *Mobile DNA*, American Society for Microbiology, Washington, D.C. 1989).

The ability of *piggyBac* to function in several dipteran species will be supportive of its use in a wider range of insects, if not other organisms. Most other vector systems function optimally, or have been only tested with their helper transposase under *hsp70* promoter regulation. The transposition efficiency of most vectors has been also found to be influenced by the amount of internal DNA inserted, the position of this DNA within the vector, and the amount of subterminal DNA remaining in the vectors.

The widespread use of *piggyBac* will be limited by the availability of easily detectable and unambiguous transformant markers. Most *Drosophila* transformations, as well as the few nondrosophilid transformations reported have depended on transformant selection by rescue of a mutant visible phenotype, usually eye pigmentation (Ashburner et al., Insect Mol. Biol., Volume 7, 201-213, 1998). Unfortunately, most insect species have neither visible mutant strains, nor the cloned DNA for the wild

type allele of the mutation, and these species require use of new dominant-acting marker genes that confer, preferably, a visible phenotype.

The present invention, discussed below, provides a system that includes vectors for transforming eukaryotic cells, derived from *piggyBac* transposons that are different from related art vectors. Furthermore, the present invention increases the transformation frequency by about eight-fold compared to other *piggyBac* transformation systems.

Summary of the Invention

It is therefore an object of the present invention to provide a transformation system contains a vector that includes DNA derived from a *piggyBac* transposon element that allows for the almost precise excision of at least a second DNA sequence that is heterologous and included in the construct and insertion of at least said second heterologous DNA sequence into eukaryotic cells after introduction of the transformation construct containing said first and at least a second DNA into said cell that is then used to form a transgenic organism wherein said transgenic organism is detectable under ultraviolet light.

Another object of the present invention is to provide a transformation system that includes a vector containing a modified *piggyBac* sequence, a sequence for marker expression linked to a polyubiquitin promoter and a nuclear localizing sequence and a helper vector including a heat shock protein gene wherein said system causes an increase in transformation frequency compared to other *piggyBac* transformation systems.

A still further object of the present invention is to provide a vector containing a modified *piggyBac* sequence and an enhanced green fluorescent protein sequence linked to a polyubiquitin promoter and a nuclear localizing sequence.

A still further object of the present invention is to provide a vector that is useful in transforming eukaryotic cells having the sequence SEQ ID No 6.

Another object of the present invention is to provide a transgenic organism that is detectable under ultraviolet light.

A further object of the present invention is to provide a eukaryotic transgenic organism that has been transformed using a transformation system that includes vector containing a modified *piggyBac* sequence, an enhanced green fluorescent protein gene linked to a polyubiquitin promoter and a nuclear localizing sequence, and a helper vector containing a heat shock protein gene promoter.

A still further object of the present invention is to provide a transgenic insect that has been transformed using a vector having the sequence SEQ ID NO 6.

Further objects and advantages of the present invention will become apparent from the following description.

Brief Description of the Drawings

Figure 1(a) is a photograph of eye color phenotypes of *Dm*[pBw] transformants.

Figure 1(b) is a photograph of a *w*[m] host strain fly (top) and orange-eye *Dm*[pBw, *gfp*] transformant fly (bottom) under brightfield (left) and UV light (right).

Figure 1 (c) is a photograph of a *w*[m] host strain fly (top) and white-eye *Dm*[pBw, *gfp*] transformant fly (bottom) under brightfield (left) and UV light (right).

Figure 2 (a) is a schematic (not to scale) of the pB[Dmw] vector showing the *Bgl*III, *Sal*I, and *Nsi*I restriction sites used to digest the genomic DNA, and the probes used for hybridization (bars). Above the schematic are distances in kilobases used to calculate internal restriction fragment sizes and minimum sizes for junction fragments. *PiggyBac* vector sequences are shaded gray, and the mini-white marker gene is white.

Figure 2 (b) shows an autoradiogram of a Southern DNA hybridization analysis of *Dm*[pBw] transformant sublines, and *w*[m] host strain control samples from transformants using the pBΔ*Sac* (experiment I) or *phsp*-pBac (experiment II) helpers using *Bgl*III digestion and *Sph*/*Hpa* *piggyBac* as probe. DNA size markers are shown to the left of the autoradiogram. M (male) and F (female)

designations refer to G0 lines, with the numbers below referring to their respective G1 sublines.

Figure 2(c) shows a Southern DNA hybridization analysis of Dm[pBw] transformant sublines and w[m] host strain control samples from transformants, using the pBASac (experiment I) or phsp-pBac (experiment II) helpers, using *SalI* digestion and *Hpa/Ase* piggyBac as probe. DNA size markers are shown to the left of the autoradiogram. M (male) and F (female) designations refer to G0 lines, with the numbers below referring to their respective G1 sublines.

Figure 2 (d) shows a Southern DNA hybridization analysis of Dm[pBw] transformant sublines and w[m] host strain control samples from transformants, using the pBASac (experiment I) or phsp-pBac (experiment II) helpers, using *NsiI* digestion and *Nsi/Hpa* + *Hpa/Nsi* probes. DNA size markers are shown to the left of the autoradiogram. M (male) and F (female) designations refer to G0 lines, with the numbers below referring to their respective G1 sublines.

Figure 3(a) is a schematic (not to scale) of the pB[Dmw, PUbnlEGFP] vector showing the *BglII*, *XhoI*, and *PstI* restriction sites used to digest the genomic DNA, and the probes used for hybridization (bars). The *Sph/Hpa* piggyBac as probe contains 0.67 kb of vector sequence (*SphI* to *BglII*) with *BglII* to *HpaI* piggyBac sequence deleted from the vector. Above the schematic are distances in kilobases used to calculate internal restriction fragment sizes and minimum sizes for junction fragments. PiggyBac vector sequences are shaded gray, the mini-white marker gene is white, and the EGFP marker gene is hatched.

Figure 3(b) is an autoradiogram of a Southern DNA hybridization analysis of Dm[pBw, *gfp*] transformant sublines, and wild type (wt) and w[m] host strain control samples using *BglII* digestion and *Sph/Hpa* piggyBac as probe. DNA size markers are shown to the left of the autoradiogram. M (male) and F (female) designations refer to G0 lines with selected G1 transformant progeny of samples.

Figure 3 (c) is an autoradiogram of a Southern DNA hybridization analysis of Dm[pBw, gfp] transformant sublines, and wild type (wt) and w[m] host strain control samples using XhoI digestion and Hpa/Ase piggyBac fragment as probe. DNA size markers are shown to the left of the autoradiogram. M and F designations refer to G0 lines with selected G1 transformant progeny of samples.

Figure 3 (d) is an autoradiogram of a Southern DNA hybridization analysis of Dm[pBw, gfp] transformant sublines, and wild type (wt) and w[m] host strain control samples using PstI digestion and Hpa/Ase piggyBac fragment + EGFP DNA as probe. DNA size markers are shown to the left of the autoradiogram. M and F designations refer to G0 lines with specific G1 line numbers are given below, with the designation (+) for those expressing visible eye pigmentation and (-) for those having non-pigmented white eyes.

Figures 4 (a) and (b) show inverse PCR strategy to isolate the pB[Dmw] vector insertion site in transformant sublines. Figure 4 (a) is a schematic (not to scale) of the vector insertion in the host plasmid showing the approximate location of the restriction sites and primers used for PCR. Forward (F) and reverse (R) primers are numbered according to their nucleotide position in piggyBac. The piggyBac sequence is shown in gray surrounded by the TTAA (SEQ ID NO 1) duplicated insertion site, the mini-white marker gene is white, and chromosomal sequence is hatched.

Figure 4(b) shows the piggyBac insertion site sequence in p3E1.2 (SEQ ID NOs 7 and 8), and the proximal insertion site sequences (SEQ ID NOs 9 and 10, 11 and 12, and 13 and 14) for three of the transformant sublines.

Figure 5 shows a circular map of the vector pB[PUB-nls-EGFP] #257.

Figures 6a-6f show SEQ ID NO 6 for pB[PUB-nls-EGFP] #257.

Figure 7(a) is a photomicrograph showing GFP expression in *Anastrepha suspensa* transformed with piggyBac/PUB-nls-EGFP at embryo stages. Under UV light, transformants exhibit bright green

fluorescence, with wild-type non-transformants exhibiting muted yellow autofluorescence (digital images taken with Leica MZ-12 fluorescence microscope and SPOT-1 CCD camera).

Figures 7(b) and 7(c) are photomicrographs showing GFP expression in *Anastrepha suspensa* transformed with *piggyBac*/PUB-nls-EGFP at larval stages. 7(b) is a wild-type non-transformant and 7c is a transformant. Under UV light, transformants exhibit bright green fluorescence, with wild-type non-transformants exhibiting muted yellow autofluorescence (digital images taken with Leica MZ-12 fluorescence microscope and SPOT-1 CCD camera).

Figures 7(d) and 7(e) are photomicrographs showing GFP expression in *Anastrepha suspensa* transformed with *piggyBac*/PUB-nls-EGFP at adult stages. 7(d) is a wild-type non-transformant and 7(e) is a transformant. Under UV light, transformants exhibit bright green fluorescence, with wild-type non-transformants exhibiting muted yellow autofluorescence (digital images taken with Leica MZ-12 fluorescence microscope and SPOT-1 CCD camera).

Figures 8(a)-8(e) are eye color phenotypes of *Bactrocera dorsalis* wild-type (+) and white eye (WE) host strain and the Bd[pBCcw] transformant lines 61, 115, and 137.

Figures 9(a) and 9(b) show medfly, *Ceratitus capitata* transformed with *piggyBac*/white/EGFP vector (pB[Ccw,pUB-nls-EGFP]) expressing eye color under brightfield (9a) and GFP expression under ultraviolet (9b).

Figure 10 shows a transgenic insect having three integrations observed under ultraviolet light after various times after decapitation. Flies were decapitated at day 0, taped in a plastic box placed outdoors in partial sunlight. Digital photographs were taken each day at the same exposure and magnification.

Detailed Description of the Invention

The present invention is an effective transformation system for producing transgenic organisms, especially transgenic insects. The identification and isolation of an autonomous *piggyBac* transposon enables transformation of cells and the production of transgenic organisms wherein DNA capable of being expressed in the

transformed cell or transgenic organism is excised from a transformation construct and inserted into the genome of a cell used to produce a transgenic organism (United States Patent Application 08/844,274; herein incorporated by reference). The term cell for the purposes of this invention includes any cell capable of being transformed by the transformation construct of the present invention and preferably includes any eukaryotic cell.

The term organism for the purposes of the present invention includes any unicellular or multicellular living entity capable of being transformed by the transformation construct of the present invention and preferably includes multicellular eukaryotes. More preferably, the cell or organism is an insect cell or an insect.

The present invention utilizes the transposon machinery of the TTAA (SEQ ID NO 1) specific transposons to excise and insert a targeted functional heterologous DNA sequence into the genome of the host cell. The resulting transformed cell or group of cells are stable transformants that are then used to make a transgenic organism, using techniques known to the skilled artisan, that will pass the introduced gene to all subsequent progeny. The targeted functional heterologous DNA for purposes of this invention is any heterologous DNA capable of being expressed in a host cell and/or a transgenic organism.

The transformation system of the present invention includes a vector, such as, for example, pB[PUB-nls-EGFP] (Figures 5 and 6), that includes a modified *piggyBac* transposon (pB), a marker construct that includes the enhanced green fluorescent protein gene (EGFP) linked to the promoter region of the *Drosophila melanogaster* polyubiquitin (PUB) gene and the nuclear localizing sequence (nls) of the SV40 virus. This vector can be used to transform and detect transgenic organisms based on expression of the green fluorescent protein marker under ultraviolet light. After chromosomal integration and inheritance of the vector, expression of green fluorescent protein occurs in all cell types, is intense, strongly localized to nuclei, and continues to be detectable under ultraviolet light even after death of the organism. The novel features of this vector includes its construction that deletes about 748 bp of internal *piggyBac*

sequence without diminishing its function, and the function of the polyubiquitin promoter in a nondrosophilid species. This has utility as a broadly based method for the creation and selection of transgenic organisms, and as a genetic marker for detecting and tracking transgenic insects used in field release programs (Figure 5).

GFP expressivity is critical for nondrosophilid species not amenable to mutant-rescue, it also widens the possibility for using the dominant expression of GFP as a primary transformant marker in many *Drosophila* lines not already carrying the *white* or *rosy* mutations, or for screens requiring selection in early development. Though vectors carrying *white* and *gfp* have been tested previously, the transformations used only *white* as the transformant selection, with GFP assessed secondarily for specific spatial or developmental expression (Davis et al., *Devel. Biol.*, Volume 170, 726-729, 1995; Wang & Hazelrigg, *Nature*, Volume 369, 400-403, 1994).

The transformation system of the present invention also includes a *piggyBac* transposase helper plasmid, pBΔSac, having its 5' terminus deleted as described by Handler et al. (1998, *supra*; herein incorporated by reference). A new transposase helper under heat-shock promoter regulation was created by the isolation of the 457 bp *Xba*I-*Xmn*I 5' nontranslated sequence from the *hsp70* gene (Lis et al., *Cell*, Volume 35, 403-410, 1983, herein incorporated by reference). The heat-shock regulated helper increases the transformation frequency by eight-fold in *Drosophila*, indicating that the *piggyBac* system could be as effective as routinely used systems such as *P* and *hobo* that have been thus far inactive in nondrosophilids (O'Brochta & Atkinson, *Insect Biochem. Molec. Biol.*, Volume 26, 739-753, 1996).

The creation of a transformed cell requires that the vector containing the functional heterologous DNA first be physically placed within the host cell. Current transformation procedures utilize a variety of techniques to introduce DNA into a cell. In one form of transformation for vertebrate systems, the DNA is microinjected directly into embryos through the use of

micropipettes. Alternatively, high velocity biolistics can be used to propel small DNA associated particles into the cell. In another form, the cell is permeablized by the presence of polyethylene glycol, thus allowing DNA to enter the cell through diffusion. DNA can also be introduced into a cell by fusing protoplasts with other entities that contain DNA. These entities include minicells, cells, lysosomes, or other fusible lipid-surfaced bodies. Electroporation is also an accepted method for introducing DNA into a cell. In this technique, cells are subject to electrical impulses of high field strength that reversibly permeabilizes biomembranes, allowing the entry of exogenous DNA sequences. One preferred method of introducing the transformation system of the present invention into insect embryos, in accordance with the present invention, is to microinject fertilized eggs with the vectors of the present invention. The DNA sequence flanked by the transposon inverted repeats will be inserted into the genome of some of the germ cells of the fertilized egg during development of the organism. This DNA will then be passed on to all of the progeny cells to produce transgenic organisms. The microinjection of eggs to produce transgenic animals has been previously described and utilized to produce transformed mammals and insects (Rubin et al., Science, Volume 218, 384-393, 1982; Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Plainview, N.Y., 1986; Morgan et al., Annu. Rev. Biochem., Volume 62, 191-217, 1993; Spradling, A.C., In: Drosophila: A Practical Approach, ed. D.B. Roberts, Oxford: IRL Press, 175-197, 1986; all herein incorporated by reference).

Accordingly, a method of producing stably transformed insects includes the step of microinjecting the transformation constructs of the present invention comprising the inverted repeats of a TTAA specific transposon and a helper construct into a cell, preferably a fertile insect egg. This is followed by incubation in an oxygenated and humidified tissue culture chamber at about 22-23° C for about 3-6 hours. Injected cells or eggs are then heat shocked at about 37°-41° C, about 39°C preferred, for about 1 hour. The resulting transformed cells or transgenic organisms have exogenous DNA inserted into the genomic DNA at the sequence TTAA.

Transformed cells and/or transgenic organisms can be selected

from untransformed cells and/or non-transgenic organisms by ultraviolet light since the transformation system includes an enhanced green fluorescent protein gene that produces an altered visible phenotype under ultraviolet light. Using standard techniques known to those familiar with the field, techniques such as, for example, Southern blotting and polymerase chain reaction, DNA can be isolated from transformed cells and/or transgenic insects to confirm that the introduced DNA has been inserted.

Genetic modification of insects with new genetic elements provides a means to control populations of agriculturally pestiferous or beneficial insects. The ability to control pest insects through genetically based sterile insect programs or genetically introduced targeted conditional susceptibilities will result in significant cost savings to agribusiness. This technology can also be used for detection and monitoring of insect populations and infestations where *piggyBac* transgenic insects are present in the population. In addition, introduction of genes that impart resistance to chemicals (including herbicides, pesticides, and insecticides) can improve the efficacy of beneficial insects. Each of these applications will result in more efficient pest control programs.

Enhancing the resistance of beneficial insects to pesticides will enhance the efficacy of the beneficial insects and may allow for the simultaneous use of chemical control and biological control of pests. Some of the beneficial insects that would make good candidates for such transformations include Hymenopteran parasitoids of *Heliothis* spp.: *Micropilitis croceipes* and *Cardiochiles nigriceps*; Hymenopteran parasitoid of Diamondback moth, *Plutella xylostella*: *Diadegma insolare*; Hymenopteran parasitoid of the Indianmeal moth, *Plodia interpunctella*: *Bracon hebitor*; and Hemipteran predators: *Xylocoris flavipes*, *Podisus maculatus*.

The following examples are intended only to further illustrate the invention and are not intended to limit the scope of the invention as described by the claims. *Drosophila melanogaster* white strain w[m], was used in the following examples

as a model system for transformation system studies using the vectors of the present invention. *D. melanogaster* and transformant progeny were maintained at about 23-25° C on standard cornmeal-yeast-molasses media.

EXAMPLE 1

The *piggyBac* transposase helper plasmid, pBASac, having its 5' terminus deleted was described previously (Handler et al., 1998; *supra*, herein incorporated by reference). pBASac was created by digestion of p3E1.2 (United States Patent Application 08/844,274) with *SacI* and religation, that deletes the 5' *piggyBac* terminal sequences but maintains the putative *piggyBac* promoter region. A transposase helper under heat-shock promoter regulation was created by isolation of the 457 bp *XbaI-XmnI* 5' nontranslated sequence from the *hsp70* gene (Lis et al., 1983, *supra*; herein incorporated by reference). The *XbaI-XmnI* fragment was blunted and ligated into the *SacI*-blunted site of pBASac to create phsp-pBac. This places the *hsp70* promoter sequence upstream of the putative *piggyBac* promoter.

The pB[Dmw] vector was created by insertion of a *Drosophila melanogaster* mini-white gene (Pirrotta et al., EMBO J., Volume 4, 3501-3508, 1985; herein incorporated by reference) into the 3E1 *piggyBac* element within the 6.0 kb p3E1.2 plasmid (Cary et al., 1989, *supra*). The mini-white gene was isolated as a 4.2 kb *EcoRI* fragment, blunted and ligated into the p3E1.2 *HpaI* site. The inserted *w* gene interrupts the *piggyBac* open reading frame (ORF), but otherwise leaves the *piggyBac* element intact, with the respective promoters in opposite orientation. A *piggyBac* vector marked with *w* and *gfp* was created by initial construction of *piggyBac* marked with an enhanced *gfp* regulated by *D. melanogaster* polyubiquitin (PUB) promoter (Lee et al., Mol. Cell. Biol., Volume 8, 4727-4735, 1988; herein incorporated by reference) linked in-frame to the SV40 nuclear localizing sequence (nls) (Lanford et al., Mol. Cell. Biol., Volume 8, 2722-2729, 1986). The

polyubiquitin-nls (PUB-nls) cassette from PUBnlsGFP (Davis et al., 1995, *supra*) was isolated as KpnI-SmaI fragment and inserted into the KpnI-SmaI cloning site of EGFP-1 (Clontech) (Cormack et al., *Gene*, Volume 173, 33-38, 1996; Yang et al., *Nucleic Acid Res.*, Volume 24, 4592-4593, 1996). Polyubiquitin-nls-EGFP was then isolated as a 4.1 kb BglII-StuI fragment and ligated into the BglII-HpaI site of piggyBac within p3E1.2 to create pB[PUBnlsEGFP]. The BglII-HpaI digestion results in a 748 bp deletion within p3E1.2. The mini-white gene was then inserted into the unique BglII site by blunt-end cloning to create pB[Dmw, PUBnlsEGFP].

EXAMPLE 2

Embryo injections used standard procedures (Rubin & Spradling, *Science*, Volume 218, 348-353, 1982; herein incorporated by reference) with dechoriation achieved either manually or by 1.6% hypochlorite solution followed by about 2 washings in approximately 0.02% Triton-X 100 in water. Eggs were placed on double-stick tape, desiccated in room-air for about 10-15 minutes and submerged under Halocarbon 700 oil. Injections followed standard *Drosophila* microinjection procedures (Rubin and Spradling, *Science*, Volume 218, 348-353, 1982; herein incorporated by reference). DNA mixtures had vector:helper concentrations of about 600:400 µg/ml, respectively, in injection buffer (approximately 5 mM KCl; approximately 0.1mM sodium phosphate; at about pH 6.8). Injected eggs were placed in an oxygenated and humidified tissue culture chamber at about 22-23°C for about 3-6 hours, and phsp-pBac injected eggs were heat shocked at about 37°C for about one hour. Hatched larvae were collected about 1-2 days later and placed on larval diet. Eclosed G0 male adults were mated either individually to about 2 or 3 w[m] virgin female adults, or in groups of about three females to about six males. G1 eggs were collected for two weeks and reared under standard conditions that include maintaining the eggs at about 23-25° C on standard cornmeal-yeast-molasses media (Ashburner et al., *supra*).

Green fluorescent protein (GFP) was observed at all

developmental stages under a Leica MZ-12 stereozoom microscope using a mercury lamp and a ultraviolet longpass filter set (HQ 41012 FITC; Chroma) optimized for red-shifted GFP variants. Photographic documentation used an Olympus OM-4 camera and 400 ASA Fujichrome film with exposure times that were determined empirically.

In the first of three transformation experiments, the piggyBac vector system was tested in *D. melanogaster* white strain using a helper transposase under piggyBac regulation (pBΔSac) and a vector marked solely with *D. melanogaster* mini-white gene (pB[Dmw]). A mixture of vector and helper plasmids at concentrations of about 600 and about 400 μg/ml, respectively, was injected into about 2,650 embryos from that about 418 larvae hatched with about 283 emerging as adults. (See Table 1 below). The G0 adults were backcrossed to w[m] flies in groups totaling about 111. Four of the G0 lines yielded G1 offspring having varying levels of eye pigmentation (Figure 1). One line (F30) was sterile, and one line produced only white eye offspring, and therefore only two of the putative Dm[pBw] transformants were verified. One of these (F13) exhibited eye pigmentation only in females in several succeeding generations, suggesting that the integration caused a sex-linked lethal mutation. Presuming a fertility rate of about 50% (fertility rates are typically between about 40-60%; see below), an approximate transformation frequency of about 1-3% of fertile G0s was obtained.

In a second experiment, the pB[Dmw] vector was again tested but with a piggyBac transposase helper under *D. melanogaster* hsp70 (Lis et al., 1983 *supra*) promoter regulation (phsp-pBac). A vector/helper mixture, at a concentration of approximately 600/400 μg/ml was injected into about 1,940 embryos, of which about 247 larvae hatched, with about 122 emerging as adults (See Table 1, below). G0 adults were initially backcrossed in a total of about 49 groups to w[m] flies, after which they were individually mated to determine fertility. Of the about 98 surviving G0 flies, about 41 yielded offspring resulting in a fertility rate of about 42%. Of the 41 fertile G0 flies, 11 lines produced offspring having

varying levels of eye coloration (Figure 1) yielding a transformation frequency of about 26%. The number of G1 offspring from the G0 lines varied considerably, ranging from 1 G1 in lines M11 and F1, to 102 G1 flies in line M13.

In a third experiment, the phsp-pBac helper was used, but with a piggyBac vector including the enhanced green fluorescent protein (gfp) marker gene in addition to the *D. melanogaster* white gene. This allowed the testing of a new gfp marker construct in transformants that could be primarily identified by white expression. Although expression of wild type GFP under polyubiquitin-nuclear localizing sequence regulation had been tested previously in *D. melanogaster* P transformants (Davis et al., 1995, *supra*), the vector of the present invention improves expression of GFP by using an enhanced GFP (EGFP-1) having a double mutation causing a reported increase in expression of up to about 35-fold (Cormack et al., 1996, *supra*; Yang et al., 1996, *supra*). The variant form is also optimized for mammalian codon usage and polyadenylation, and preliminary tests of the marker construct indicated transient GFP expression in both *Drosophila* embryos and dipteran and lepidopteran cell lines (A.M. Handler and R.A. Harrell, unpublished). The vector construct, pB[Dmw, PUBnlEGFP], also allowed evaluation of piggyBac transformation with about a 10.0 kb vector, approximately 3.4 kb larger than previous vectors tested, and having about 748 bp of piggyBac DNA deleted (previous vectors retained all piggyBac DNA). As before, a mixture of about 600 µg/ml vector and about 400 µg/ml helper was injected into about 2147 embryos, of which about 412 larvae hatched, and about 218 emerged as adults (Table 1 below). G0 adults were backcrossed to w[m] flies in a total of about 90 mating groups, of which about 79 yielded offspring. Although white gene expression (eye pigmentation) was depended upon as the primary marker, G1 larvae and pupae were examined under UV for visible GFP expression, and seven of the G0 lines yielded fluorescent G1 larvae and pupae. Interestingly, as shown below in Table 2, upon adult emergence only six of the seven G0 lines

yielded G1 offspring with observable eye color pigmentation. While about 70 G1 offspring in total exhibited observable green fluorescence, only about 27 of these flies exhibited a level of eye pigmentation that would have allowed their selection under normal screening procedures. In contrast, all of the G1 flies with eye color pigmentation expressed GFP. Figure 1b shows a *Dm[pBw, egfp]* transformant having an orange eye color and GFP fluorescence, with no fluorescence observed in the *w[m]* host. Figure 1c shows another transformant having a white eye phenotype indistinguishable from that in the *w[m]* host strain, but exhibiting an equal, if not greater level of GFP fluorescence compared to the orange eye transformant. Notably, fluorescence is quenched in the eye of the pigmented transformant, while it is easily visible in the white eye transformant. High magnification examination revealed a few pigmented ommatidia in some white eye G1 flies expressing GFP, though these would not have been normally detected. Based on selection by GFP expression and presuming about 50% fertility, an approximate transformation frequency of about 6-7% of fertile G0 flies is deduced.

An assessment of vector activity based on germline transformation frequency is a factor of both transposon mobility in the host embryo and levels of genomic position effect suppression of the marker gene, or stated more simply, the ability to visibly identify putative transformants. While position effect variegation and suppression of white expression in transformants is well established (Hazelrigg et al., *Cell*, Volume 64, 1083-1092, 1984; Pirodda et al., 1985, *supra*), the effect of complete marker suppression on transformation frequencies has not been assessed since such transformants have been only detected fortuitously after molecular analysis. The experiment using both the white and GFP markers proved the importance of position effects on marker expression convincingly, since GFP was readily detectable in 70 G1 flies, yet eye pigmentation was apparent in less than 40% of these. Under typical screening procedures these flies would not have been scored as transformants, though pigmentation in a few ommatidia in some flies could be detected at high magnification, and for a few lines, pigmentation was more apparent in subsequent

generations. It is likely that expression of the white marker would have been improved by heat shock regulation, but nonetheless, GFP was easily detected in all the non-pigmented transformants, and strongly expressed in some. The influence of modifier genes on position effect variegation is complex, and target genes (or their promoters) are not equivalently affected (Bhadra et al., Genetics, Volume 150, 251-263, 1998). The polyubiquitin-gfp gene may be a target of position effect modifiers, but it is clearly less susceptible to suppression relative to white in terms of its expressed phenotype in the same chromosomal context. The data suggests that GFP is a more reliable visible marker than white, that portends well for its use as a general marker in other insects.

TABLE 1. Transformation Experiments.

Expt	vector/ helper	eggs injected	G0s mated	% fertility	No. G0 lines	No. G1 lines	transformant frequency
I	pB[Dmw]/pBΔSac	2,650	283	nd	4	11	0.01-0.03*
II	pB[Dmw]/phsp-pBac	1,940	122	42	11	266	0.26
III	pB[Dmw, PUbnlEGFP]/phsp- pBac	2,147	218	nd	7	70	0.06-0.07*

*estimated frequency based on 50% fertility

TABLE 2. G1 white⁺ and GPR marker expression in Dm[pBw, gfp] transformants.

G0 line	No. G1	GFP	white ⁺	Frequency white ⁺
M4	4	4	3	0.75
M9	21	21	2	0.10
M12	3	3	1	0.33
M23	15	15	14	0.93
M45	5	5	0	0
M47	21	21	6	0.29
F10	1	1	1	1.00
Total	70	70	27	0.39

EXAMPLE III

Southern hybridization was performed to verify genomic transposition of the *piggyBac* vectors. Approximately 5-10 µg of genomic DNA was digested with indicated restriction enzymes and separated on about 0.8% agarose gels. DNA was stained with ethidium bromide, blotted to nylon filters and immobilized by ultraviolet irradiation. Hybridization probes were labeled with [³²P]-dCTP by random priming (Gibco BRL) according to the manufacturer's specifications. Probe DNA was generated from indicated *piggyBac* restriction fragments (see below) that were separated from p3E1.2, or the entire *egfp* gene from pEGFP-1 (Clontech) by agarose electrophoresis and gel-elution. Hybridizations were performed in phosphate buffer, approximately pH 7.5; about 1% BSA; about 7% SDS at about 65°C with an initial wash in about 2X SSC; about 0.2% SDS at about room temperature and about two washes in about 1X SSC; about 0.1% SDS at about 55°C for approximately 30 minutes. Autoradiography was performed by exposure of Kodak X-Omat film at about -90° C.

Genomic transposition of the *piggyBac* vectors was verified by Southern DNA hybridization. The basic strategy was to perform hybridizations to the 5' vector arm using the *piggyBac* *SphI*-*HpaI* or *NsiI*-*HpaI* fragment as probe, and the 3' vector arm using the *HpaI*-*AseI* or *HpaI*-*NsiI* fragment as probe. Using probes to both

vector arms, internal fragments spanning most of the vector were detected. Hybridizations to the vector arms and adjacent chromosomal sequence indicate their presence in non-plasmid DNA and indicate the number of integrations, while internal hybridizations that yield known fragment sizes confirm vector integrity.

For pB[Dmw] transformants, genomic DNA was initially digested with *Bgl*III and hybridized to the labeled *Sph*-*Hpa* piggyBac fragment, that detects both vector arms resulting in two bands for each integration (Figure 2A). Each intact vector integration should result in one band greater than about 0.67 kb for the 5' arm, and one band greater than about 5.9 kb for the 3' arm. Since varying eye color phenotypes among G1 sublines was observed, and in some cases within G1 sublines, sublines having light orange, dark orange, or red eye coloration from the same G1 sublines were selected for hybridization analysis. For example, flies having differing phenotypes from lines M13-39, M19-90, and M19-91 were hybridized separately, but no difference in the number or sites of insertion were apparent. Of all the lines tested, all had single integrations except for two lines having two integrations (M13-39 and M19-91) and one line having three integrations (F14-63). All the lines with multiple integrations had dark orange or red eye color, though several lines with a single integration also shared these phenotypes. Hybridization patterns for the lines tested indicated that for most of the G0 lines, different integrations were transmitted to many of the G1 sibling offspring. For example, the three G1 sublines tested from both the M3 and M5 G0 lines all show different patterns indicating at least three independent integrations occurring in the two G0 germlines.

Genomic DNA digested with *Sal*I and hybridized to *Hpa*I-*Ase*I probe yielded single bands greater than about 3.0 kb for each integration, and the number of integrations determined were consistent with the *Sph*I-*Hpa*I hybridizations (Figure 2B). For all samples, *Nsi*I digestion and hybridization to *Nsi*-*Hpa*I and *Hpa*I-*Nsi*I probe yielded only about 1.5 kb and about 4.6 kb bands accounting for about 6.1 kb of the about 6.6 kb vector, indicating the same generally high level of vector integrity for all

integrations tested.

G1 sublines from six G0 lines transformed with the pB[Dmw, PUBnlEGFP] vector were digested with either *Bgl*II and probed with *Sph*I-*Hpa*I piggyBac DNA for 5' vector arm analysis, or digested with *Xho*I and probed with *Hpa*I-*Ase*I piggyBac DNA for 3' arm analysis (Figure 3A and 3B). Both hybridizations yielded one band for each sample, indicating single integrations having occurred in each line. *Nsi*I restriction digests with *Nsi*I-*Hpa*I and *Hpa*I-*Nsi*I hybridizations yielded about 0.7 kb and about 0.8 kb bands indicating vector integrity for each integration (data not shown).

Two G0 lines, M9 and M47, yielded a high proportion of G1 flies expressing only GFP and white eyes, and line M45 that yielded only white eye transformants. These lines were analyzed by *Pst*I digestion and hybridization to EGFP and *Hpa*-*Ase*. All lines shared the about 4.4 kb internal vector fragment, with an additional junction fragment from the 3' vector arm and adjacent insertion site chromosomal DNA. The M9 white eye lines all shared the same integration indicated by an about 0.9 kb junction fragment, and similarly the M47 white eye lines all shared the same 5.0 kb junction fragment. The pigmented lines M9-2 and M9-3 had different integrations from each other, and from their white eye sibling lines, and the pigmented lines M47-9 and M47-10 shared the same integration based on an about 4.0 kb junction fragment, but which differs from their white eye siblings. These hybridizations, and that for M45-1, proves that the white eye flies were transformed, and that white expression was likely influenced by differing insertion sites from their pigmented sibling lines.

EXAMPLE IV

To verify that piggyBac-mediated chromosomal transpositions had occurred, insertion sites were isolated by inverse PCR from sublines F1-2, M17-4 and M31-6, all having single integrations. Inverse PCR was performed as described previously (Handler et al., 1998, *supra*; herein incorporated by reference) using *Hae*III digestions for 5' and 3' junctions and *Msp*I digestion for 3'

junctions. After about 4 hours digestion, restriction fragments were circularized by ligation at about 16° C for about 16 hours. PCR was performed on the circularized fragments by using primer sequences in opposite orientation within the *piggyBac* restriction site and terminus for each junction. For the 5' junction, the forward primer (572F) 5'-TCTTGACCTTGCCACAGAGG-3' (SEQ ID NO 2) and reverse primer (154R) 5'-TGACACTTACCGCATTGACA-3' (SEQ ID NO 3) were used. For the 3' junction the reverse primer (2118R) 5'-GTCAGTCCAGAAACAACCTTTGGC-3' (SEQ ID NO 4) and the forward primer (2385F) 5'-CCTCGATATACAGACCGATAAAAACACATG-3' (SEQ ID NO 5) were used. PCR products were separated in low-melting-temperature agarose, and fragments were selected that were longer than the respective restriction site terminus distances and different from those expected from the p3E1.2 based vector and helper plasmids. These products were directly subcloned into ddT vectors (Invitrogen), that were sequenced by using primers to vector sequence proximal to the respective termini. Subcloned PCR products were sequenced and analyzed by alignment using GeneWorks 2.5 software (Oxford Molecular Group) and subjected to BLAST analysis (Altschul et al., J. Mol. Biol., Volume 215, 403-410, 1990; herein incorporated by reference) to identify genomic insertion site sequences and distinguish them from those in the injected plasmids. For all the integrations both the 5' and 3' junctions yielded the *piggyBac* inverted terminal repeat sequences immediately adjacent to a TTAA sequence (SEQ ID NO 1) and proximal insertion site DNA (Figure 4). The TTAA (SEQ ID NO 1) duplicated target site is characteristic of all *piggyBac* integrations (Elick et al., Genetica, Volume 97, 127-139, 1995) and typically indicates a vector-mediated transposition. The BLAST analysis revealed that the M17-4 integration occurred in a TTAA site within the *cubitus interruptus*-Dominant gene located on chromosome 4 at nucleotide 12,898 (GenBank submission U66884; Ahmed & Podemski, Gene, Volume 197, 367-373, 1997), and the M3106 integration was found to have occurred in a TTAA site within a previously sequenced region of the distal X chromosome (GenBank submission AL09193; Murphy et al, direct submission). Determination of insertions in these previously sequenced sites gives the first

direct proof that a *piggyBac* vector does indeed insert into and duplicates TTAA (SEQ ID NO 1) insertion sites in a eukaryotic genome.

Two of the three insertion sites that were sequenced were found to be in previously sequenced genomic loci, and as expected, the insertion sites were all TTAA (SEQ ID NO 1) with one of them within the *ci^D* allele on the fourth chromosome. Many transposons have insertion site preferences, and for at least some, a clear negative bias against specific sites or loci. This has been clearly demonstrated by genomic hotspots and coldspots for *P* integration in *D. melanogaster* (See Engels, In: *Mobile DNA*, D.E. Berg and M.M. Howe, eds., American Society of Microbiology, Washington, D.C., 439-484, 1989), and by differences in preferential integration sites between *hobo* and *P* (Smith et al., *Genetics*, Volume 135, 1063-1076, 1993). If the TTAA (SEQ ID NO 1) specificity for *piggyBac* integration is not further influenced by proximal sequences, then *piggyBac* transpositions may find use in transposon-mutagenesis and enhancer traps for loci refractory to *P* or *hobo* transpositions in *Drosophila*.

Example V

The Caribbean fruit fly, *Anastrepha suspensa*, was transformed with a *piggyBac* vector marked solely with PUB-nls-GFP (pB[PUB-nls-EGFP]) (Figures 5 and 6) using the *hsp70-piggyBac* (phsp-pBac) helper. From injected embryos, 561 surviving G0 adults were intermated in 60 small groups. Four of the G0 groups yielded a total of 57 G1 offspring exhibiting green fluorescence at all stages of development (See Figure 7) and chromosomal vector integrations were verified by Southern hybridization for each G0 group. To test GFP as a genetic marker for field released transgenic flies, the perdurance of GFP expression was assayed in transgenic flies killed by decapitation. Two to three day old *A. suspensa* adults transformed with pB[PUB-nls-EGFP], and wild type non-transformed adults, were decapitated and placed within a plastic box kept outdoors in partial shade. GFP fluorescence was

observed daily by digital images taken with a SPOT-1 cooled CCD digital camera (Diagnostic Instruments, Inc.) through a Leica MZ-12 stereozoom microscope. All images were taken at the same magnification and exposure parameters. Figure 10 shows that while GFP fluorescence decreases with time after death, unambiguous detection of GFP is still possible at 28 days after decapitation, with no fluorescence detectable in wild flies. This indicates that the PUB-nls-EGFP marker should be a reliable visible detection system for released transgenic insects, and especially for those captured and killed in field traps with monitoring occurring after extended time periods.

Example VI

A *piggyBac* vector marked with the Mediterranean fruit fly (*Ceratitidis capitata*) white gene cDNA (pB[Ccw]) and the phsp-pBac helper was used to transform the oriental fruit fly (*Bactrocera dorsalis*). Injected G0 embryos from the *B. dorsalis* white eye mutant strain yielded 102 fertile adults, that upon individual backcrossing, yielded three lines of putative transformants with pigmented eyes (Figures 8a-8e). One of these lines produced 119 G1 transformants. Southern DNA hybridization analysis with *piggyBac* and white gene probe verified chromosomal integration of the *piggyBac*-white vector in all three lines. In a separate experiment, the white/PUB-nls-EGFP marker within pB[Ccw, PUB-nls-EGFP] was introduced into a single *B. dorsalis* transformant line from 17 G0 matings. As in *Drosophila*, the transformant was selected solely by GFP expression, having undetectable eye coloration. This reaffirms the notion that the polyubiquitin-EGFP marker is significantly more reliable than white gene markers.

Example VII

The PUB-nls-EGFP marker was introduced into the medfly, *Ceratitidis capitata*, to further test GFP as a transgenic selection, and to create GFP-marked strains for testing as a field release marker in medfly SIT. First a *piggyBac* vector marked with PUB-

nls-GFP and the medfly *white* gene (pB[Ccw, PUb-nls-EGFP]) was tested, and then the vector solely marked with Pub-nls-GFP pB[PUb-nls-EGFP]) was tested. Both experiments used the *hsp70-piggyBac* (phsp-pBAC) helper. Based on GFP fluorescence, the first experiment yielded five transformant lines from 99 fertile G0s (See Figure 9), while the second experiment yielded three transformed lines from 17 fertile G0s. Transformation was verified by Southern hybridization analysis. The foregoing detailed description is for the purpose of illustration. Such detail is solely for that purpose and those skilled in the art can make variations without departing from the spirit and scope of the invention.

We claim:

1. A transformation system comprising a vector containing a nucleotide sequence from a piggyBac transposon in which is inserted an enhanced green fluorescent protein gene linked to a promoter region of a polyubiquitin gene and a nuclear localizing sequence of an SV40 virus.
2. A transformation system of claim 1 further comprising a piggyBac transposase helper plasmid under heat-shock promoter regulation.
3. The transformation system of claim 1 wherein said polyubiquitin gene is from *Drosophila melanogaster*.
4. The transformation system of claim 2 wherein said polyubiquitin gene is from *Drosophila melanogaster*.
5. The transformation system of claim 1 wherein said piggyBac transposon is modified by deleting about 748 bp of internal piggyBac sequence.
6. A vector having SEQ ID NO 6.
7. A transgenic organism transformed using the transformation system of claim 3 wherein in said organism is detectable under ultraviolet light.
8. A transgenic insect transformed using the transformation system of claim 3 wherein said insect is detectable under ultraviolet light.
9. The transgenic insect of claim 8 that continues to be detectable under ultraviolet light after death.



FIG. 1a



FIG. 1b



FIG. 1c




Fig. 2a

*Bgl*II digestion - *Sph*/Hpa probe




FIG. 2b

*Sa*II digestion - *Hpa*/*Ase* probe




FIG. 2c

*Nsi*I digestion - *Nsi*I/*Hpa* + *Hpa*/*Nsi* probes




FIG. 2d




Fig. 3a

*Bgl*II digestion - *Sph*/*Hpa* probe *Xho*I digestion - *Hpa*/*Ase* probe

kb M9 M4 M12 M23 M47 F10 M9 M4 M12 M23 M47 F10 wt




FIG. 3b FIG. 3c

*Pst*I digestion - *Hpa*/*Ase*+EGFP probe




FIG. 3d




Fig. 4a

p3E1.2 aagcgcaaatcttttTTAA-piggyBac-TTAAataatagtttcttaat

F1-2 aaaagactgactatTTAA-piggyBac-TTAAaagcacactgagtc

M17-4 aaatgtcgtctaggTTAA-piggyBac-TTAAagccgtatatcagat

M31-6 aaatgaacgacttttTTAA-piggyBac-TTAAtggttttttagttgt

FIG. 4b




FIG. 5

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
GACGAAAGGG	CCTCGTGATA	CGCCTATTTT	TATAGGTTAA	TGTCATGATA	50
ATAATGGTTT	CTTAGACGTC	AGGIGGCACT	TTTCGGGGAA	ATGIGCGCGG	100
AACCCCTATT	TGTTTATTTT	TCTAAATACA	TTCAAATATG	TATCCGCTCA	150
TGAGACAATA	ACCTTGATAA	ATGCTTCAAT	AATATTGAAA	AAGGAAGAGT	200
ATGAGTATTC	AACATTTCCG	TGTGCGCCTT	ATTCCCTTTT	TTCGGGCATT	250
TTGCTTCCCT	GTTTTTGCTC	ACCCAGAAAC	GCTGGTGAAA	GTAAAAGATG	300
CIGAAGATCA	GTGGGGTGCA	CGAGTGGGTT	ACATCGAACT	GGATCTCAAC	350
AGCGGTAAAG	TCCTTGAGAG	TTTTCGCCCC	GAAGAAGGTT	TTCCAATGAT	400
GAGCACITTT	AAAGTTCIGC	TATGTGCGCC	GGTATTATCC	CGTATTGACG	450
CCGGGCAAGA	GCAACTCCGT	CGCCGCATAC	ACTATTCTCA	GAATGACTTG	500
GTGAGTACT	CACCAGTCAC	AGAAAAGCAT	CTTACGGATG	GCATGACAGT	550
AAGAGAATTA	TGCAGTCTTG	CCATAACCAT	GAGTGATAAC	ACTGCGGCCA	600
ACTTACTTCT	GACAACGATC	GGAGGACCGA	AGGAGCTAAC	CGCTTTTTTG	650
CACAACATGG	GGGATCATGT	AACTCGCCTT	GATCGTTGGG	AACCGGAGCT	700
GAATGAAGCC	ATACCAAACG	ACGAGCGTGA	CACCAAGATG	CCGTGTAGCA	750
TGGCAACAAC	GTTCGCGCAA	CTATTAACTG	GCGAACTACT	TACTCTAGCT	800
TCCCGGCAAC	AATTAAATAG	CTGGATGGAG	GCGGATAAAG	TTGCAGGACC	850
ACTTCTGGCC	TCCGCGCCTT	CGGCTGGCTG	GTTTATTGCT	GATAAATCTG	900
GAGCCGGTGA	GCGTGGGTCT	CGCGGTATCA	TTGCAGCACT	GGGGCCAGAT	950
GGTAAGGCGT	CCCGTATCGT	AGTTATCTAC	ACGACGGGGA	GTCAGGCAAC	1000
TATGGATGAA	CGAAATAGAC	AGATCGCTGA	GATAGGTTGC	TCACTGATTA	1050
AGCATTTGGT	ACTGTTCAGC	CAAGTTTACT	CATATATACT	TTAGATTGAT	1100
TTAAAACTTC	ATTTTAAATT	TAAAAGGATC	TAGGTGAAGA	TCCTTTTTGA	1150
TAATCTCATG	ACCAAAATCC	CTTAACGTGA	GTTTTGTTTC	CACGTAGCGT	1200
CAGACCCCGT	AGAAAAGATC	AAAGGATCTT	CTTGAGATCC	TTTTTTTCTG	1250
CGCGTAATCT	GCTGCTTGCA	AACAAAAAAA	CCACCGCTAC	CAGCGGTGGT	1300
TTGTTTGGCG	GATCAAGAGC	TACCAACTCT	TTTTCCGAAG	GTAACTGGCT	1350
TCAGCAGAGC	GCAGATACCA	AATACTGTCC	TTCTAGTGTA	GCGGTAGTTA	1400
GGCCACCCTT	TCAAGAACTC	TGTAGCACCG	CCTACATAAC	TGGCTCTGCT	1450
AATCCCTGTA	CCAGTGGCTG	CTGCCAGTGG	CGATAAGTCC	TGCTTTACCG	1500
GGTTGGACTC	AAGACGATAG	TTACCGGATA	AGGCGCAGCG	GTGGGGCTGA	1550
ACGGGGGGTT	CGTGCACACA	GCCCAGCTTG	GAGCGAACGA	CCTACACCGA	1600
ACTGAGATAC	CTACAGCGTG	AGCATTGAGA	AAGCGCCACG	CTTCCCGAAG	1650
GGAGAAAGGC	GGACAGGTAT	CCGGTAAGCG	GCAGGGTCCG	AACAGGAGAG	1700
CGCACGAGGG	AGCTTCCAGG	GGGAAACGCC	TGGTATCTTT	ATAGTCCCTG	1750
CGGGTTTCGC	CACCTCTGAC	TTGAGCGTCC	ATTTTGTGTA	TGCTGTGACG	1800

FIG. 6a

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
GGGGGOGGAG	CCTATGGAAA	AACGCCAGCA	ACGGGGCCTT	TTTACGGTTC	1850
CTGGCCTTTT	GCIGGCCTTT	TGCICACATG	TTCCTTCTG	CGTTATCCCC	1900
TGATTCTGTG	GATAACGGTA	TTACCGCCTT	TGAGIGAGCT	GATACCGCTC	1950
GGCGCAGCCG	AACGACCGAG	CGCAGCGAGT	CAGIGAGCGA	GGAGCGGAA	2000
GAGCGCCCAA	TACGCAAACC	GCTCTCCCC	GCGCGTTGGC	CGATTCAATTA	2050
ATGCAGCTGG	CACGACAGGT	TTCGGACIG	GAAAGCGGGC	AGIGAGCGCA	2100
AOGCAATTAA	TGIGAGTTAG	CTCACTCATT	AGGCACCCCA	GGCTTTACAC	2150
TTTATGCTTC	CGGCTCGTAT	GTGTGTGTGA	ATTGTGAGCG	GATAACAATT	2200
TCACACAGGA	AACAGCTATG	ACCATGATTA	CGAATTCGAG	CTCGGTACCC	2250
GGGGATCCTC	TAGAGTCGAC	CTGCAGGCAT	GCAAGCTTGC	ATGCTGCGAG	2300
GTGGAAGCTC	GCGGACTTGG	GTTCGCCATT	CTTTAGCGCG	CGTCGGGICA	2350
CACAGCTTGG	CCACAATGTG	GTTTTGTGCA	AACGAAGATT	CTATGACGTG	2400
TTTAAAGTTT	AGGTGAGTAA	AAGCGCAAT	CTTTTTHAAC	CCTAGAAAGA	2450
TAGTCTGGGT	AAAATTGACG	CATGCATTCT	TGAAATATTG	CTCTCTCTTT	2500
CTAAATAGCG	CGAATCCGTC	GCTGTGCATT	TAGGACATCT	CAGTCGCGGC	2550
TTGGAGCTCC	CGTGAGGCGT	GCTGTGCAAT	GCGGTAAGTG	TCACTGATTT	2600
TGAACATATA	CGACCGCGTG	AGTCAAAATG	ACGCATGATT	ATCTTTTACG	2650
TGACTTTTAA	GATTTAACCT	ATACGATAAT	TATATTGTTA	TTTCATGTTC	2700
TACTTAAGTG	ATACTTATT	ATATATATAT	TTCTTGTTA	TAGATATCGT	2750
GACTAATATA	TAATAAAATG	GGTAGTCTCT	TAGACGATGA	GCATATCCTC	2800
TCCTGCTCTC	TGCAAGCGGA	TGACGAGCTT	GTGGGIGAGG	ATTCTGACAG	2850
TGAAATATCA	GATCAAGTAA	GIGAAGATGA	CGTCAGAGC	GATACAGAAG	2900
AAGCGTTTAT	AGATGAGGTA	CATGAAGTGC	AGCCAACGTC	AAGCGGTAGT	2950
GAAATATTAG	ACGAACAAA	TGTTATTGAA	CAACAGGTT	CTTCATTGGC	3000
TTCTAACAGA	ATCTTGACCT	TGCCACAGAG	GACTATTAGA	GGTAAGAATA	3050
AACATTGTGT	GTCAACTTCA	AAGTCCAGCA	GGGTAGCCG	AGTCTCTGCA	3100
CTGAACATTG	TCAGATCTCG	AGCTCAAGCT	TGGAATCTG	CAGTCGACCG	3150
TACCGGATCT	TGTGCGCGGA	ACGCAGCGAC	AGAGATTCCA	ATGTGTCCGT	3200
ATCTTTTACG	CTTTTGCCCT	TCAGTTCAG	ACGAAGCGAC	TGGCGATTCT	3250
CGTGTGGGGT	CTGCTTCAGG	GCTTGTGAA	TTAGGGGCGG	CAGATCGCCG	3300
ATGGGGGTGG	CGCGGAGGG	CACTTTCACC	TTCGCGTACG	GCTTGTCTGT	3350
CTTCGGGTTC	AAAATCTCCA	GCTCCATTTT	GCTTTGGGIG	CGCTTGCAAT	3400
CAGTACTGTG	CAAAATCGAA	AATCGCCGAA	CCGTAGTGTG	ACCGTGGGG	3450
GCTCTGCGAA	AATAAACCTT	TTTAGGTATA	TGCCACACA	CGGGGAAGC	3500
ACAGTGGATT	ATATGTTTTA	ATATTATAAT	ATGCAGGTTT	TCATTACTTA	3550
TCCAGATGTA	AGCCCACTTA	AAGCGATTTA	ACAATTATTT	GCCGAAAGAG	3600

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
TAAAAACAAA	TTTCACTTAA	AAATGGATTA	AGAAAAGCTT	GIGTAAGATT	3650
ATGCGCAGCG	TTGCCAGATA	GCTCCATTTA	AAACACTTCA	AAAACAATAA	3700
GT'TTIGAAAA	TATATACATA	AATAGCAGTC	GT'TGCGGCAA	CGCTCAACAC	3750
ATCACACTTT	TAAACACCC	TTTACCTACA	CAGAATTACT	TTTTAAATTT	3800
CCAGTCAAGC	TGCGAGTTTC	AAAATTATAG	CCGGTAGAGA	AGACAGIGCT	3850
ATTTCAAAAG	CAACTAAAT	AAACACCAAT	CCTAACAAGC	CTTGGACTTT	3900
TGTAAGTTTA	GATCAAAGGT	GGCATTGCAT	TCAATGTCAT	GGTAAGAAGT	3950
AGGTGCTCTA	GGTAGAAATC	CTCATTCAGC	CGGTCAAGTC	AGTAAGAGAA	4000
AGGTCTCAAT	TTGAAATTGT	CTTAAAAATA	TTTTATTGTT	TGTACTGTG	4050
GTTGAGTTTA	ACGAAAAACA	CAAAAAAAA	GIGATACACA	GAAATCATAA	4100
AAAATTTTAA	TACAAGGTAT	TGGTACGTAT	CAAAACATT	TGGGCACAAT	4150
TTTTTTTCTC	TGTACTAAAG	TGTTACGAAC	ACTACGGTAT	TTTTTAGTGA	4200
TTTTTCAACG	ACACCGAAGG	TATATAAACA	GCGTTGCGGA	ACGGTGGCCT	4250
TCAAAAOCAA	TTGACATTIG	CAGCAGCAAG	TACAAGCAGA	AAGTAAGCG	4300
CAATCAGCGA	AAAATTTATA	CTTAATTGTT	CGTGATTAAA	GTACAATTAA	4350
AAGAACATTC	TGGAAAGTCA	CAAGAAACGT	AAGTTTTTAA	CTGGCTGTTA	4400
OCAATTAGTA	ATAAGAGCAA	CAAGACGTIG	AGTAATTICA	AGAAAACTIG	4450
CATTTCAAGG	TCTTTGTTGG	GCCATTTTTT	TTTTATTCAA	CGCTCTACGT	4500
AATTACAAAA	TAAGAAATTG	GCAGCCACGC	ATCTTGTTTT	CCCAATCAAT	4550
TGGCATCAAA	ACGCAACAA	ATCTATAAAT	AAAACITGCG	TGTTGATTTT	4600
CGCCAAGATT	TATTGGCAAA	TGTGAAATT	CGCAGTGAAG	CATTTGAAAA	4650
TTCGAGAAAT	CAOGAACCA	CTCGAGCATT	TGTGTGCAATG	TTATTAGTTA	4700
GT'TAGTTCTT	TGCTTAATTG	AAGTATTTTA	OCAACGAAT	CCACTTATTT	4750
TLAGCTGAAA	TAGAGTAGGT	TGCTTGAAC	GAAAGCCACG	TCGTGAAAT	4800
TTCTTATTGC	TTAGTAGTTG	TGACGTACCC	ATATACACAC	AAAATAATGT	4850
GLATGCATGC	GTTTCAGCTG	TGTATATATA	CATGCACACA	CTGGCATTAT	4900
GAAAAAGATG	ACGAGCAACG	GAACAGGTTT	CTCAACTACC	TTTGTTCCTG	4950
TT'TCTTGGCT	TTCCTTTGTT	CCAATATTGG	TAGAGGGTTA	ATAGGGGTTT	5000
CTCAACAAAG	TTGGCGTGG	TAAATAAGTT	TCCATTTTTT	ATTCCCCAGC	5050
CAGGAAGTTA	GTTTCAATAG	TTTTGTAAAT	TCAACGAAC	TCATTIGATT	5100
TGGTACTAAT	TTTCCACATC	TCTATTTTGA	CCCGCAGAAT	AATCCAAAT	5150
GCAGATCGGG	GATCCACACC	CACCCAAGAA	GAAGCGCAAG	GTTGAGGACG	5200
ATCCCGTGGT	TTTACAACGT	CGTGACTGGG	AAAACCTGG	CGTTACCCAA	5250
CT'TAATGGCC	TTCAGCACA	TCCCCCTTTC	GCCAGCTGGC	GTAATAGCGA	5300
AGAGGCCCCG	ACCGATGGCC	CTTCCCAACA	GTTCGGGTGG	ACTCTAGAGG	5350
ATCCCCGGGA	TCCACGGTTC	GCCACCATGG	TGAGCAAGGG	CGAGGAGCTG	5400

FIG. 6 c

14/22

SUBSTITUTE SHEET (RULE 26)

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
TTCACCGGGG	TGGTGGCCAT	CCTGGTGGAG	CTGGACGGGG	ACGTAAACGG	5450
CCACAAGTTC	AGCGTGTCCG	GCGAGGGCGA	GGGGATGGCC	ACCTACGGCA	5500
AGCTGACCCCT	GAAGTTCATC	TGCACCAACG	GCAAGCTGGC	CGTGGCCCTGG	5550
CCCAACCCCTG	TGACCAACCCCT	GACCTACGGC	GTCAGTGTCT	TCAGCCGCTA	5600
CCCGACCCAC	ATGAAGCAGC	ACGACTTCTT	CAAGTCCGGC	ATGCCCCGAG	5650
GCTACGTCCA	GGAGCGCACC	ATCTTCTTCA	AGGAAGACGG	CAACTACAAG	5700
ACCCGGGGCG	AGGTGAAGTT	CGAGGGGGAC	ACCCGTGGTA	ACCGCATCGA	5750
GCTGAAGGGC	ATCGACTTCA	AGGAGGACGG	CAACATCCIG	GGGCACAAGC	5800
TGGAGTACAA	CTACAACAGC	CACAACGTCT	ATATCATGGC	CGACAAGCAG	5850
AAGAACGGCA	TCAAGGIGAA	CTTCAAGATC	CGCCACAACA	TCGAGGACGG	5900
CAGCGTGCAG	CTGGCCGACC	ACTACCAGCA	GAACACCCCC	ATCGGGGACG	5950
GGCCCGTGTCT	GCTGCCCCGAC	AACCACTACC	TGAGCACCCA	GTCGGCCCTG	6000
AGCAAAGACC	CCAACGAGAA	GGCGATCAC	ATGGTCTGTC	TGGAGTTCTG	6050
GACCGCCGGC	GGGATCACTC	TCGGCATGGA	CGAGCTGTAC	AAGTAAAGGG	6100
GGCGCGACTC	TAGATCATAA	TCAGCCATAC	CACATTGTGA	GAGGTTTAC	6150
TTGCTTTTAA	AAACCTCCCA	CACCTCCCCC	TGAACCTGAA	ACATAAAATG	6200
AATGCAATTG	TTGTGTGTAA	CTGTGTTATT	GCAGCTTATA	ATGGTTACAA	6250
ATAAAGCAAT	AGCATCACAA	ATTTACAAA	TAAAGCATTT	TTTTCACTGC	6300
ATTCTAGTTG	TGGTTGTGCC	AAACATCA	ATGTATCTTA	AGGCGTAAAT	6350
TGTAAGCGTT	AATATTTTGT	TAAATTCGC	GTTAAATTTT	TGTTAAATCA	6400
GCTCATTTTT	TAACCAATAG	GGCGAAATCG	GCAAAATCCC	TTATAAATCA	6450
AAAGAATAGA	CCGAGATAGG	GTTGAGTGT	GTTCCAGTTT	GGAACAAGAG	6500
TCCACTATTA	AAGAAGTGG	ACTCCAAAGT	CAAAGGGCGA	AAAACCGICT	6550
ATCAGGGCGA	TGGCCCACTA	CGTGAACCAT	CACCTAATC	AAGTTTTTTG	6600
GGGTCCAGGT	GGGTAAAGC	ACTAAATCGG	AACCTTAAAG	GGAGCCCCCG	6650
ATTTAGAGCT	TGACGGGGAA	AGCCGGCGAA	CGTGGCGAGA	AAGGAAGGGA	6700
AGAAAGCGAA	AGGAGCGGGC	GCTAGGGGCG	TGGCAAGTGT	AGCGGTACCG	6750
CTGCGCGTAA	CCACCACACC	CGCCGGCTT	AATGGCGCGC	TACAGGGGCG	6800
GTCAGGTGGC	ACTTTTCGGG	GAAATGTGGG	CGGAACCCCT	ATTTGTTTAT	6850
TTTTCTAAAT	ACATTCAAAT	ATGTATCCGC	TCATGAGACA	ATAACCCCTG	6900
TAAATGCTTC	AATAATATTG	AAAAAGGAAG	AGTCCGTAGG	CGGAAGAAGC	6950
CAGCTGTGGA	ATGTGTGTCA	GTTAGGGTGT	GGAAAGTCCC	CAGGCTCCCC	7000
AGCAGGCAGA	AGTATGCAAA	GCATGCATCT	CAATTAGTCA	GCAACCAGGT	7050
GTTGAAAGTC	CCAGGCTCC	CCAGCAGGCA	GAAGTATGCA	AAGCATGCAT	7100
CTCAATTAGT	CAGCAACCAT	AGTCCCGGCC	CTAACTCCGC	CCATCCCGCC	7150
CCTAACCTCC	CCAGTTCCG	CCATTCTCC	GGCCCATGGC	TGACTAATTT	7200

FIG. 6d

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
TTTTTATTTA	TGCAGAGGOC	GAGGCGGCT	CGGOCICIGA	GCTATTCCAG	7250
AAGTAGIGAG	GAGGCTTTTT	TGCAGGAACC	ATTGIGGGAA	COGTGCGATC	7300
AAACAAACGC	GAGATACCGG	AAGTACIGAA	AAACAGIUGC	TCCAGGOCAG	7350
TGGGAACATC	GATGTTTTGT	TTTGACGGAC	CCCTTACTCT	CGTCTCATAT	7400
AAACCGAAGC	CAGCTAAGAT	GGTATACCTA	TTATCATCTT	GIGATGAGGA	7450
TGCTTCTATC	AACGAAAGTA	CCGGTAAACC	GCAATCGGTT	ATGTATTATA	7500
ATCAAACATA	AGGCGGAGTG	GACACGCTAG	ACCAAATGIG	TTCGTGIGATG	7550
AOCITGCAGTA	GGAAGACGAA	TAGGTGGGCT	ATGGCATTTAT	TGTACCGAAT	7600
GATAAACATT	CCCTGCATAA	ATTCITTTTAT	TATATACAGC	CATAATGICA	7650
GTAGCAAGGG	AGAAAAGGTC	CAAAGTCGCA	AAAAATTTAT	GAGAAACCTT	7700
TACATGAGCC	TGACGTCATC	GTTTATGCGT	AAGCGTTTAG	AAGCTCCTAC	7750
TTTGAAGAGA	TATTTGCGCG	ATAATATCTC	TAATATTTTG	CCAAATGAAG	7800
TGOCITGGTAC	ATCAGATGAC	AGTACTGAAG	AGCCAGTAAT	GAAAAAACGT	7850
ACTTACTGTA	CTTACTGCCC	CTCTAAAATA	AGGCGAAAGG	CAAATGCATC	7900
GIGCAAAAAA	TGCAAAAAAG	TTATTGTGCG	AGAGCATAAT	ATTGATATGT	7950
GCCAAAGTTG	TTCTGACTG	ACTAATAAGT	ATAATTGTTT	TCTATTATGT	8000
ATAAGTTAAG	CTAATTACTT	ATTTTATAAT	ACAACATGAC	TGTTTTTAA	8050
GTACAAAATA	AGTTTATTTT	TGTAAAAGAG	AGAATGTTTA	AAAGTTTGT	8100
TACTTTATAG	AAGAAATTTT	GAGTTTGTGT	TTTTTTTAA	TAAATAAATA	8150
AACATAAATA	AATGTTTGT	TGAATTTATT	ATTAGTATGT	AAGTGTAAT	8200
ATAATAAAAC	TTAATATCTA	TTCAAATTAA	TAAATAAACC	TCGATATACA	8250
GACCGATAAA	ACACATGCGT	CAATTTTACG	CAATGATTATC	TTTAACTGAC	8300
GTCACATAT	GATTATCTTT	CTAGGGTTAA	ATAATAGTTT	CTAATTTTTT	8350
TATTTATCAG	CCCTGCTGCG	TGAATACCGT	ATATCTCAAC	GCCTGCTGIG	8400
AGATTGTGCGT	ATCTTAGGCT	TTTGTAGTTT	TGGCTCATUG	ACTGTATATT	8450
GTCOGACACA	TTTGTGTGGA	TTTGTGTTTT	GATCAAGAC	TTGAGCAGAG	8500
ACACGTAAAT	CAACGTGTTA	AATGTATCCA	TATTAAAGAT	ATCAACCGCA	8550
TGCGTATATG	GTCGTAAAA	TATATTTTTT	AACCTCTTA	TACTTTGCAC	8600
TCTTGGTTAA	TACGGTTGCG	TGTACAGACG	TAATCATGTT	TCTTTTTTTG	8650
GATAAAACTC	CTACTGAGTT	TGACCTCATA	TTAGACCTTC	ACAAGTTGCA	8700
AAACGTGGCA	TTTTTTTACCA	ATGAAGAAAT	TAAAGTTATT	TTAAAAAATT	8750
TCAATCACAGA	TTTAAAGAAG	AACCAAAAAT	TAAATTATTT	CAACAGTTTA	8800
ATCGACCAAT	TAATCAACGT	GTACACAGAC	GCGTCGGCAA	AAAACACGCA	8850
GCCCGACGIG	TTGGCTAAAA	TTATTAAATC	AACITGIGTT	ATAGTCACGG	8900
ATTGTGCGTC	CAACGTGTTT	CTCAAAAAGT	TGAAGACCAA	CAAGTTTACG	8950
GACACTATTA	ATTATTIGAT	TTTGCCCCAC	TTCATTTTGT	GGGATCACAA	9000

FIG. 6e

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
TTTIGTTATA	TTTAAACAA	AGCTTGGCAC	TGGCCGTGGT	TTTACAAAGT	9050
OGTGACTGGG	AAAACCCCTGG	CGTTACCCAA	CTTAATCGGC	TTCAGCACA	9100
TCCCCCTTTC	GCCAGCTGGC	GTAATAGCGA	AGAGGCCCCG	ACCGATCGGC	9150
CTTCCCAACA	GTTCGCCAGC	CTGAATGGCG	AATGGGGCCT	GATCGGGTAT	9200
TTTCTCCTTA	CGCATCTGIG	CGGTATTICA	CACCGCATAT	GGTGCACCTT	9250
CAGTACAATC	TGCTCTGATG	CCGCATAGTT	AAGCCAGCCC	CGACACCCGC	9300
CAACACCCGC	TGACGGGGCC	TGACGGGCTT	GTCCTGCTCC	GGCATCCGCT	9350
TACAGACAAG	CTGTGAACGT	CTCCGGGAGC	TGCATGTGTC	AGAGGTTTTC	9400
ACCGTCAICA	CCGAAACGGC	CGA			9423

FIG. 6 f



FIG. 7a



FIG. 7b



FIG. 7c



FIG. 7d




FIG. 7e




FIG. 8a




FIG. 8b




FIG. 8c




FIG. 8d




FIG. 8e



FIG. 9a



FIG. 9b




FIG. 10

SEQUENCE LISTING

<110> Handler, Alfred M.

<120> PiggyBac Transformation System

<130> 0194.98

<140>

<141>

<160> 14

<170> PatentIn Ver. 2.0 - beta

<210> 1

<211> 4

<212> DNA

<213> Trichoplusia ni

<400> 1

ttaa

4

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 2

tcttgacctt gccacagagg

20

<210> 3

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 3

tgacacttac cgcattgaca

20

<210> 4

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 4

gtcagtcag aaacaacttt ggc

23

<210> 5

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 5

cctcgatata cagaccgata aaaacacatg

30

<210> 6

<211> 9423

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial
Sequence:pB [PUb-nls-EGFP] #257

<400> 6

gacgaaaggg cctcgtgata cgcctatddd tataggdtaa tgcctatgata ataattggddd 60
 cttagacgtc aggtggcact tttcggggaa atgtgcgcgg aaccctatt tgtttatddd 120
 tctaaataca ttcaaatatg tatcgcgtca tgagacaata accctgataa atgcttcaat 180
 aatattgaaa aaggaagagt atgagtattc aacatttccg tgcgcgcctt attcctddd 240
 ttgcggcatt ttgccttctt gtttttgctc acccagaaac gctgggtgaaa gtaaaagatg 300
 ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac agcggtaaga 360
 tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactddd aaagttctgc 420
 tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggg cggcgcatat 480
 actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg 540
 gcatgacagt aagagaatta tgcagtgctg ccataaccat gagtataac actgcggcca 600
 acttacttct gacaacgatc ggaggaccga aggagctaac cgcttttttg cacaacatgg 660
 gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacg 720
 acgagcgtga caccacgatg cctgtagcaa tggcaacaac gttgcgcaaa ctattaactg 780
 gcgaactact tactctagct tcccggcaac aattaataga ctggatggag gcggataaag 840
 ttgcaggacc acttctgcgc tcggcccttc cggctggctg gtttattgct gataaatctg 900
 gagccggtga gcgtgggtct cgcggtatca ttgcagcact ggggccagat ggtaagccct 960

cccgtatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa cgaaatagac 1020
agatcgctga gatagggtgcc tcactgatta agcattggta actgtcagac caagtttact 1080
catatatact ttagattgat ttaaaacttc atttttaatt taaaaggatc taggtgaaga 1140
tcctttttga taatctcatg accaaaatcc cttaacgtga gttttcgttc cactgagcgt 1200
cagaccccgt agaaaagatc aaaggatctt cttgagatcc tttttttctg cgcgtaatct 1260
gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgg ttgtttgccg gatcaagagc 1320
taccaactct ttttccgaag gtaactggct tcagcagagc gcagatacca aatactgtcc 1380
ttctagtgtg gccgtagtta ggccaccact tcaagaactc tgtagcaccg cctacatacc 1440
tcgctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg 1500
ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga acgggggggtt 1560
cgtgcacaca gccagcttg gagegaacga cctacaccga actgagatac ctacagcgtg 1620
agcattgaga aagcgccacg cttcccgaag ggagaaaggc ggacaggtat ccggtaaagc 1680
gcagggtcgg aacaggagag cgcacgaggg agcttccagg gggaaacgcc tggatatctt 1740
atagtcctgt cgggtttcgc cacctctgac ttgagcgtcg atttttgtga tgctcgtcag 1800
gggggcggag cctatggaaa aacgccagca acgcggcctt tttacgggtc ctggcctttt 1860
gctggccttt tgctcacatg ttctttcctg cgttatcccc tgattctgtg gataaccgta 1920
ttaccgcctt tgagtgaact gataccgctc gccgcagccg aacgaccgag cgcagcagat 1980
cagtgaagca ggaagcggaa gagegcecaa tacgcaaacc gcctctcccc gcgcgttggc 2040
cgattcatta atgcagctgg cacgacaggt ttcccgaact gaaagcgggc agtgagcgca 2100
acgcaattaa tgtgagttag ctactcatt aggcacccca ggctttacac tttatgcttc 2160
cggctcgtat gttgtgtgga attgtgagcg gataacaatt tcacacagga aacagctatg 2220
accatgatta cgaattcgag ctcggtaccc ggggatcctc tagagtcgac ctgcaggcat 2280
gcaagcttgc atgcctgcag gtcgacgctc gcgcgacttg gtttgccatt ctttagcgcg 2340
cgtcgcgtca cacagcttgg ccacaatgtg gtttttgtca aacgaagatt ctatgacgtg 2400
tttaaagttt aggtcgagta aagcgcaaat cttttttaac cctagaaaga tagtctgcgt 2460
aaaattgacg catgcattct tgaaatattg ctctctcttt ctaaatagcg cgaatccgtc 2520
gctgtgcatt taggacatct cagtcgccgc ttggagctcc cgtgaggcgt gcttgtcaat 2580
gcggtaagtg tcactgattt tgaactataa cgaccgcgtg agtcaaatg acgcatgatt 2640
atcttttacg tgacttttaa gatttaactc ataogataat tatattgtta tttcatgttc 2700

tacttacgtg ataacttatt atatatatat tttcttgta tagatategt gactaatata 2760
taataaaatg ggtagttctt tagacgatga gcatatcctc tctgctcttc tgcaaagcga 2820
tgacgagctt gttggtgagg attctgacag tgaaatatca gatcacgtaa gtgaagatga 2880
cgtccagagc gatacagaag aagcgtttat agatgaggta catgaagtgc agccaacgtc 2940
aagcggtagt gaaatattag acgaacaaaa tgttattgaa caaccagggtt cttcattggc 3000
ttctaacaga atcttgacct tgccacagag gactattaga ggtaagaata aacattgttg 3060
gtcaacttca aagtccacga ggcgtagccg agtctctgca ctgaacattg tcagatctcg 3120
agctcaagct tcgaattctg cagtcgacgg taccgatct tgcgcgcga acgcagcgac 3180
agagattcca atgtgtccgt atctttcagg cttttgccct tcagttccag acgaagcgac 3240
tgggcattcg cgtgtggggg ctgcttcagg gtcttgtaaa ttagggcgcg cagatcgccg 3300
atgggcgtgg cgccggaggg caccttcacc ttgccgtacg gcttgctgtt cttcgcgctc 3360
aaaatctcca gctccatttt gctttcgggtg cgcttgcaat cagtactgtc caaaatcgaa 3420
aatcgccgaa ccgtagtgtg accgtgcggg gctctgcgaa aataaacttt tttaggtata 3480
tggccacaca cggggaaagc acagtggatt atatgtttta atattataat atgcaggttt 3540
tcattactta tccagatgta agcccactta aagcgattta acaattattt gccgaaagag 3600
taaaaacaaa tttcacttaa aaatggatta agaaaagctt gtgtaagatt atgcgcagcg 3660
ttgccagata gctccattta aaacacttca aaaacaataa gttttgaaaa tatatacata 3720
aatagcagtc gttgcgcgaa cgctcaacac atcacacttt taaaacaccc tttacctaca 3780
cagaattact ttttaaattt ccagtcaagc tgcgagtttc aaaattatag ccggtagaga 3840
agacagtgtt atttcaaaag caaactaaat aaacaccaat cctaacaagc cttggacttt 3900
tgtaagttta gatcaaagggt ggcattgcat tcaatgtcat ggtaagaagt aggtcgtcta 3960
ggtagaaatc ctcattcagc cgggtcaagtc agtacgagaa aggtctcaat ttgaaattgt 4020
cttaaaaata ttttattggt ttgtactgtg gtgagtttaa acgaaaaaca caaaaaaaaa 4080
gtgatacaca gaaatcataa aaaattttta tacaagggtat tcgtacgtat caaaaacatt 4140
tcggcacaat tttttttctc tgtactaaag tgttacgaac actacgggtat tttttagtga 4200
ttttcaacgg acaccgaagg tatataaaca gcgttcgcga acggtcgcct tcaaaaccaa 4260
ttgacatttg cagcagcaag tacaagcaga aagtaaagcg caatcagcga aaaatttata 4320
cttaattggt ggtgattaaa gtacaattaa aagaacattc tcgaaagtca caagaaacgt 4380

aagtttttaa ctcgctgtta ccaattagta ataagagcaa caagacgttg agtaatttca 4440
agaaaaactg catttcaagg tctttgttcg gccatttttt ttttattcaa cgctctacgt 4500
aattacaaaa taagaaattg gcagccacgc atcttgtttt cccaatcaat tggcatcaaa 4560
acgcaaacia atctataaat aaaacttgcg tgttgatttt cgccaagatt tattggcaaa 4620
ttgtgaaatt cgcagtgcg catttgaaaa ttcgagaaat cacgaacgca ctcgagcatt 4680
tgtgtgcatg ttattagtta gttagtctt tgcttaattg aagtatttta ccaacgaaat 4740
ccacttattt ttagctgaaa tagagtaggt tgcttgaaac gaaagccacg tctggaaaat 4800
ttcttattgc ttagtagttg tgacgtcacc atatacacac aaaataatgt gtatgcatgc 4860
gtttcagctg tgtatatata catgcacaca ctcgcattat gaaaacgatg acgagcaacg 4920
gaacagggtt cteaactacc tttgttcctg tttcttcgct ttcctttgtt ccaatattcg 4980
tagaggggta atagggggtt ctcaacaaag ttggcgtcga taaataagtt tcccattttt 5040
attccccagc caggaagtta gtttcaatag ttttgtaatt tcaacgaaac tcatttgatt 5100
tcgtactaat tttccacatc tctattttga cccgcagaat aatccaaaat gcagatcggg 5160
gatccccacc cacccaagaa gaagcgcaag gtggaggacg atcccgctcg tttacaacgt 5220
cgtgactggg aaaaccctgg cgttacccaa cttaatcgcc ttgcagcaca tccccctttc 5280
gccagctggc gtaatagcga agaggcccg accgatcgcc ctcccaaca gttgcggtcg 5340
actctagagg atccccggga tccaccggtc gccaccatgg tgagcaaggg cgaggagctg 5400
ttcaccgggg tgggtgccc atctggtcgag ctggacggcg acgtaaacgg ccacaagttc 5460
agcgtgtccg gcgagggcga gggcgatgcc acctacggca agctgaccct gaagttcatc 5520
tgcaccaccg gcaagctgcc cgtgccctgg cccaccctcg tgaccaccct gacctacggc 5580
gtgcagtgtc tcagccgcta ccccgaccac atgaagcagc acgacttctt caagtccgcc 5640
atgcccgaag gctacgtcca ggagcgcacc atcttcttca aggacgacgg caactacaag 5700
acccgcgccg aggtgaagtt cgagggcgac accctggtga accgcacga gctgaagggc 5760
atcgacttca aggaggacgg caacatcctg gggcacaagc tggagtacaa ctacaacagc 5820
cacaacgtct atatcatggc cgacaagcag aagaacggca tcaaggtgaa cttcaagatc 5880
cgccacaaca tcgaggacgg cagcgtgcag ctcgccgacc actaccagca gaacaccccc 5940
atcggcgacg gccccgtgct gctgcccgc aaccactacc tgagcaccca gtccgccctg 6000
agcaaagacc ccaacgagaa gcgcgatcac atggtcctgc tggagttcgt gaccgccgcc 6060
gggatcactc tcggcatgga cgagctgtac aagtaaagcg gccgcgactc tagatcataa 6120

tcagccatac cacatttgta gaggttttac ttgctttaa aaacctcca cacctcccc 6180
tgaacctgaa acataaaatg aatgcaattg ttgttgtaa cttgtttatt gcagcttata 6240
atggttacia ataaagcaat agcatcacia atttcacaaa taaagcattt ttttcaactgc 6300
attctagtgt tggtttgtcc aaactcatca atgtatctta aggcgtaaat tgtaagcgtt 6360
aatattttgt taaaattcgc gttaaatttt tgttaaata gctcattttt taaccaatag 6420
gccgaaatcg gcaaaatccc ttataaatca aaagaataga ccgagatagg gttgagtgtt 6480
gttccagttt ggaacaagag tccactatta aagaacgtgg actccaacgt caaagggcga 6540
aaaacgtct atcagggcga tggcccacta cgtgaaccat caccctaate aagttttttg 6600
gggtcgaggt gccgtaaagc actaaatcgg aaccctaaag ggagcccccg atttagagct 6660
tgacggggaa agccggcgaa cgtggcgaga aaggaaggga agaaagcgaa aggagcgggc 6720
gctagggcgc tggcaagtgt agcggtcacg ctgcgcgtaa ccaccacacc cgccgcgctt 6780
aatgcgcgc tacagggcgc gtcaggtggc acttttcggg gaaatgtgcg cggaaccctt 6840
atttgtttat ttttctaat acattcaaat atgtatccgc tcatgagaca ataaccctga 6900
taaagtcttc aataatattg aaaaaggaag agtcctgagg cggaaagaac cagctgtgga 6960
atgtgtgtca gttaggggtg ggaaagtccc caggctcccc agcaggcaga agtatgcaaa 7020
gcatgcatct caattagtca gcaaccaggt gtggaaagtc cccaggctcc ccagcaggca 7080
gaagtatgca aagcatgcat ctcaattagt cagcaaccat agtccccgcc ctaactccgc 7140
ccatcccgcc cctaactccg ccagttccg cccattctcc gccccatggc tgactaattt 7200
ttttatttta tgcagaggcc gaggcgcct cggcctctga gctattccag aagtagtgag 7260
gaggtttttt tggaggaacc attgtgggaa ccgtgcgac aaacaaacgc gagataccgg 7320
aagtactgaa aaacagtcgc tccaggccag tgggaacatc gatgttttgt tttgacggac 7380
cccttactct cgtctcatat aaacegaagc cagctaagat ggtatactta ttatcatctt 7440
gtgatgagga tgcttctatc aacgaaagta ccggtaaacc gcaaatgggt atgtattata 7500
atcaaactaa aggcggagtg gacacgctag accaaatgtg ttctgtgatg acctgcagta 7560
ggaagacgaa taggtggcct atggcattat tgtacggaat gataaacatt gcctgcataa 7620
attcttttat tatatacagc cataatgtca gtagcaaggg agaaaaggtc caaagtcgca 7680
aaaaatttat gagaaacctt tacatgagcc tgacgtcatc gtttatgcgt aagcgtttag 7740
aagctectac tttgaagaga tatttgccgcg ataatatctc taatattttg ccaaatgaag 7800

tgcctggtac atcagatgac agtactgaag agccagtaat gaaaaaacgt acttactgta 7860
cttactgccc ctctaaaata aggcgaaagg caaatgcate gtgcaaaaaa tgcaaaaaag 7920
ttatttgtcg agagcataat attgatatgt gccaaagttg tttctgactg actaataagt 7980
ataatttgtt tctattatgt ataagttaag ctaattactt attttataat acaacatgac 8040
tgtttttaaa gtacaaaata agtttatttt tgtaaaagag agaagtgtta aaagttttgt 8100
tactttatag aagaaatttt gagtttttgt ttttttttaa taaataaata aacataaata 8160
aattgtttgt tgaatttatt attagtatgt aagtgtaaat ataataaaac ttaatatcta 8220
ttcaaattaa taaataaacc tcgatataca gaccgataaa acacatgcgt caattttacg 8280
catgattatc tttaacgtac gtcacaatat gattatcttt ctaggggttaa ataatagttt 8340
ctaatttttt tattattcag cctgctgtcg tgaataccgt atatctcaac gctgtctgtg 8400
agattgtcgt attctagcct ttttagtttt tcgctcatcg acttgatatt gtccgacaca 8460
tttctgtcga tttgcgtttt gatcaaagac ttgagcagag acacgttaat caactgttca 8520
aattgatcca tattaacgat atcaaccoga tgcgtatatg gtgcgtaaaa tatatttttt 8580
aaccctctta tactttgcac tctgcgttaa taecgcttcg tgtacagacg taatcatgtt 8640
ttcttttttg gataaaactc ctactgagtt tgacctcata ttagaccctc acaagttgca 8700
aaacgtggca ttttttacca atgaagaatt taaagttatt ttaaaaaatt tcatcacaga 8760
tttaaagaag aaccaaaaat taaattattt caacagttta atcgaccagt taatcaacgt 8820
gtacacagac gcgtcggcaa aaaacacgca gcccgacgtg ttggctaaaa ttattaaatc 8880
aacttgtgtt atagtcacgg atttgccgtc caacgtgttc ctcaaaaagt tgaagaccaa 8940
caagtttacg gacactatta attatttgat tttgccccac ttcattttgt gggatcacaa 9000
ttttgttata ttttaaacaa agcttggcac tggccgtcgt tttacaacgt cgtgactggg 9060
aaaaccctgg cgttacccaa cttaatcgcc ttgcagcaca tcccccttc gccagctggc 9120
gtaatagcga agaggcccc accgatcgcc cttcccaaca gttgcgcagc ctgaatggcg 9180
aatggcgct gatgcggtat tttctctta cgcactctgt cggtatttca caccgcatat 9240
ggtgcactct cagtacaatc tgctctgatg ccgcatagtt aagccagccc cgacacccgc 9300
caacacccgc tgacgcgccc tgacgggctt gtctgctccc ggcacccgct tacagacaag 9360
ctgtgaccgt ctccgggagc tgcattgtgc agaggtttt accgtcatca ccgaaacgcg 9420
cga 9423

<210> 7
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:p3E1.2

<400> 7
aagcgcaaatt cttttttaa

19

<210> 8
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:p3E1.2

<400> 8
ttaaataata gtttctaatt

19

<210> 9
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:F1-2

<400> 9
aaaaagactg actatttaa

19

<210> 10
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:F1-2

<400> 10
ttaataagca cactgagtc

19

<210> 11
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:M17-4

<400> 11

aaaatgtcgt ctaggttaa

19

<210> 12

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:M17-4

<400> 12

ttaaagccgt atatcagat

19

<210> 13

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:M31-6

<400> 13

aaatgaacga cttttttaa

19

<210> 14

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:M31-6

<400> 14

ttaatgggtt ttttagttgt

19

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/22433

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 15/00, 15/63; A01K 67/00, 67/027

US CL : 435/320.1, 455; 800/13, 14, 19, 20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 455; 800/13, 14, 19, 20

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, STN, MEDLINE, CAPLUS, BIOSIS, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HANDLER et al., The Lepidopteran Transposon Vector, piggybac, Mediates germ-Line Transformation in the Mediterranean Fruit Fly. Proc. Natl. Acad. Sci., USA. June 1998, Vol. 95, pp. 7520-7525, especially pages 7520, 7521.	1-9
Y	CORMACK et al. ACS-Optimized Mutants of the Green Fluorescent Protein (GFP). Science. 1996, Vol. 173, pp. 33-38, especially pages, 33, 38.	1-9
Y	DAVIS et al. A Nuclear GFP That Marks Nuclei in Living Drosophila Embryos; Maternal Supply Overcomes a Delay in the Appearance of Zygotic Fluorescence. Developmental Biology. 1995, Vol. 170, pp. 726-729, especially pages, 726, 729.	1-9

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
I document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 22 SEPTEMBER 2000	Date of mailing of the international search report 02 NOV 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Shin-Lin Chen</i> SHIN-LIN CHEN Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/22433

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	O'BROCHTA et al. Transposable Elements and Gene Transformation in Non-Drosophilid Insects, Insect Biochem. Molec. Biol., 1996, Vol. 26, Nos. 8-9, pp. 739-753, especially pages 739, 740, 749, 750.	7-9
A	ASHBURNER et al. Prospects for the Genetic Transformation of Arthropods. Insect Molecular Biology. 1998, Vol. 7, No. 3, pp. 201-213, especially pages 201, 205.	7-9

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.